


SUPPLEMENTARY MATERIAL TO
**On the dynamics of some small structural motifs in
 rRNA upon ligand binding**

ALEKSANDRA RAKIĆ^{1#} and PETAR M. MITRAŠINOVIĆ^{2*#}

¹Faculty of Physical Chemistry, University of Belgrade, Studentski Trg 12-16, 11000
 Belgrade and ²Center for Multidisciplinary Studies, University of Belgrade,
 Kneza Višeslava 1, 11030 Belgrade, Serbia

J. Serb. Chem. Soc. **73** (1) (2008) 41–53

Secondary structures generated by the RNADRAW program¹ for 21- and 27-nu-
 cleotide sequences in the immediate vicinity of the maximum- $\Delta(\eta, \theta)$ ligand bin-
 ding sites denoted by A2581, A1354 and A764 in the 1njn, 1nwy and 1jzx com-
 plexes, respectively:



#Serbian Chemical Society member.

*Corresponding author. E-mail: petar.mitrasinovic@cms.bg.ac.yu

A1354

5'

3'

[illegible]

27-nucleotide sequence

1. I. L. Hofacker, W. Fontana, P. F. Stadler, L. S. Bonhoeffer, M. Tacker, P. Schuster, *Monats. Chem.* **125** (1994) 167.